arXiv:1101.0075v2 [math.CA] 4 Jan 2011

ON THE SIMPSON'S INEQUALITY FOR CO-ORDINATED CONVEX FUNCTIONS

M. EMIN ÖZDEMIR[♦], AHMET OCAK AKDEMIR[♠,★], HAVVA KAVURMACI[♦],
AND MERVE AVCI[♦]

Abstract. In this paper, a new lemma is proved and inequalities of Simpson type are established for co-ordinated convex functions and bounded functions.

1. INTRODUCTION

The following inequality is well-known in the literature as Simpson's inequality:

Theorem 1. *Let* $f:[a,b]\rightarrow\mathbb{R}$ *be a four times continuously differentiable mapping on* $[a,b]$ *and* $\left\Vert f^{(4)}\right\Vert_{\infty}=\sup\limits_{x\in[a,b]}\left\vert f^{(4)}(x)\right\vert<\infty$*. Then the folllowing inequality holds:*

$$\left\vert \frac{1}{3}\left[\frac{f(a)+f(b)}{2}+2f\left(\frac{a+b}{2}\right)\right]-\frac{1}{b-a}\int_{a}^{b}f(x)\,dx\right\vert \leq \frac{1}{2880}\left\Vert f^{(4)}\right\Vert_{\infty}(b-a)^{4}.$$

For recent results on Simpson's type inequalities see the papers [12], [13], [14], [15], [16], [17], [18] and [19].

In [11], Dragomir defined co-ordinated convexity and proved the following inequalities:

Let us consider the bidimensional interval $\Delta=[a,b]\times[c,d]$ in $\mathbb{R}^{2}$ with $a<b$ and $c<d$. A function $f:\Delta\rightarrow\mathbb{R}$ will be called convex on the co-ordinates if the partial mappings

$$f_{y}:[a,b]\rightarrow\mathbb{R},\quad f_{y}(u)=f(u,y)$$

and

$$f_{x}:[c,d]\rightarrow\mathbb{R},\quad f_{x}(v)=f(x,v)$$

are convex where defined for all $y\in[c,d]$ and $x\in[a,b]$.

Recall that the mapping $f:\Delta\rightarrow\mathbb{R}$ is convex on Δ, if the following inequality;

$$f(\lambda x+(1-\lambda)z,\lambda y+(1-\lambda)w)\leq\lambda f(x,y)+(1-\lambda)f(z,w)$$

holds for all $(x,y),(z,w)\in\Delta$ and $\lambda\in[0,1]$.

Date: December 29, 2010.

2000 *Mathematics Subject Classification.* 26D10,26D15.

Key words and phrases. Simpson's inequality, co-ordinates, convex functions .

★Corresponding Author.

Theorem 2. *Suppose that* $f:\Delta=[a,b]\times[c,d]\rightarrow\mathbb{R}$ *is convex on the co-ordinates on* Δ*. Then one has the inequalities;*

$$\begin{aligned}
& f\left(\frac{a+b}{2},\frac{c+d}{2}\right) \\
& \leq \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx \qquad (1.1)\\
& \leq \frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{4}
\end{aligned}$$

The above inequalities are sharp.

Recently, several papers have been written on co-ordinated convexity. In [8], Hwang et al. gave a refinement of Hadamard's inequality on the co-ordinates and they proved some inequalities for co-ordinated convex functions. In [5], [6] and [7], Alomari and Darus proved inequalities for co-ordinated $s-$convex functions. In [1], Akdemir and Özdemir gave definition of co-ordinated $P-$convex functions and Godunova-Levin functions and proved some inequalities. In [2], Latif and Alomari defined co-ordinated $h-$convex functions and established some inequalities for co-ordinated $h-$convex functions. In [3], Latif and Alomari proved inequalities involving product of convex functions on the co-ordinates. In [9], Özdemir et al. defined co-ordinated $m-$convexity and $(\alpha,m)-$convexity and gave inequalities. In [10], Sarıkaya et al. proved a new lemma and established some inequalities on co-ordinated convex functions. In [4], Bakula and Pecaric gave several Jensen-type inequalities for co-ordinated convex functions.

In this paper, we give a Simpson-type inequality for co-ordinated convex functions on the basis of the following lemma.

2. MAIN RESULTS

To prove our main result, we need the following lemma.

Lemma 1. *Let* $f:\Delta\subset\mathbb{R}^2\rightarrow\mathbb{R}$ *be a partial differentiable mapping on* $\Delta=[a,b]\times[c,d]$*. If* $\frac{\partial^2 f}{\partial t\partial s}\in L(\Delta)$*, then the following equality holds:*

$$\begin{aligned}
& \frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9} \\
& +\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36} \qquad (2.1)\\
& -\frac{1}{6(b-a)}\int_a^b\left[f(x,c)+4f\left(x,\frac{c+d}{2}\right)+f(x,d)\right]dx \\
& -\frac{1}{6(d-c)}\int_c^d\left[f(a,y)+4f\left(\frac{a+b}{2},y\right)+f(b,y)\right]dy \\
& +\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx \\
= \ & (b-a)(d-c)\int_0^1\int_0^1 p(x,t)\,q(y,s)\,\frac{\partial^2 f}{\partial t\partial s}(ta+(1-t)b,sc+(1-s)d)\,dtds
\end{aligned}$$

where

$$p\left(x,t\right)=\begin{cases}\left(t-\frac{1}{6}\right), & t\in\left[0,\frac{1}{2}\right]\\ \left(t-\frac{5}{6}\right), & t\in\left(\frac{1}{2},1\right]\end{cases}$$

and

$$q\left(y,s\right)=\begin{cases}\left(s-\frac{1}{6}\right), & s\in\left[0,\frac{1}{2}\right]\\ \left(s-\frac{5}{6}\right), & s\in\left(\frac{1}{2},1\right]\end{cases}.$$

Proof. Integrating by parts, we can write

$$\begin{aligned}&\int_0^1\int_0^1 p\left(x,t\right)q\left(y,s\right)\frac{\partial^2 f}{\partial t\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)dtds\\ =\;&\int_0^1 q\left(y,s\right)\left[\int_0^{\frac{1}{2}}\left(t-\frac{1}{6}\right)\frac{\partial^2 f}{\partial t\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)dt\right.\\ &\left.+\int_{\frac{1}{2}}^1\left(t-\frac{5}{6}\right)\frac{\partial^2 f}{\partial t\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)dt\right]ds\end{aligned}$$

By integrating the right hand side of equality, we get

$$\begin{aligned}&\int_0^1 q\left(y,s\right)\left\{\left[\left(t-\frac{1}{6}\right)\left(\frac{1}{a-b}\right)\frac{\partial f}{\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)\right]_0^{\frac{1}{2}}\right.\\ &-\frac{1}{a-b}\int_0^{\frac{1}{2}}\frac{\partial f}{\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)dt\\ &+\left[\left(t-\frac{5}{6}\right)\left(\frac{1}{a-b}\right)\frac{\partial f}{\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)\right]_{\frac{1}{2}}^1\\ &\left.-\frac{1}{a-b}\int_{\frac{1}{2}}^1\frac{\partial f}{\partial s}\left(ta+\left(1-t\right)b,sc+\left(1-s\right)d\right)dt\right\}ds\end{aligned}$$

$$
\begin{aligned}
= \quad & \frac{1}{b-a}\Bigg\{-\frac{1}{3}\int_0^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\frac{\partial f}{\partial s}\left(\frac{a+b}{2},sc+(1-s)\,d\right)ds \\
& -\frac{1}{3}\int_{\frac{1}{2}}^1\left(s-\frac{5}{6}\right)\frac{\partial f}{\partial s}\left(\frac{a+b}{2},sc+(1-s)\,d\right)ds \\
& -\frac{1}{6}\int_0^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\frac{\partial f}{\partial s}\left(b,sc+(1-s)\,d\right)ds \\
& -\frac{1}{6}\int_{\frac{1}{2}}^1\left(s-\frac{5}{6}\right)\frac{\partial f}{\partial s}\left(b,sc+(1-s)\,d\right)ds \\
& +\int_0^{\frac{1}{2}}\int_0^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\frac{\partial f}{\partial s}\left(ta+(1-t)\,b,sc+(1-s)\,d\right)dsdt \\
& +\int_{\frac{1}{2}}^1\int_0^{\frac{1}{2}}\left(s-\frac{5}{6}\right)\frac{\partial f}{\partial s}\left(ta+(1-t)\,b,sc+(1-s)\,d\right)dsdt \\
& -\frac{1}{6}\int_0^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\frac{\partial f}{\partial s}\left(a,sc+(1-s)\,d\right)ds \\
& -\frac{1}{6}\int_{\frac{1}{2}}^1\left(s-\frac{5}{6}\right)\frac{\partial f}{\partial s}\left(a,sc+(1-s)\,d\right)ds \\
& -\frac{1}{3}\int_0^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\frac{\partial f}{\partial s}\left(\frac{a+b}{2},sc+(1-s)\,d\right)ds \\
& -\frac{1}{3}\int_{\frac{1}{2}}^1\left(s-\frac{5}{6}\right)\frac{\partial f}{\partial s}\left(\frac{a+b}{2},sc+(1-s)\,d\right)ds \\
& +\int_0^{\frac{1}{2}}\int_{\frac{1}{2}}^1\left(s-\frac{1}{6}\right)\frac{\partial f}{\partial s}\left(ta+(1-t)\,b,sc+(1-s)\,d\right)dsdt \\
& +\int_{\frac{1}{2}}^1\int_{\frac{1}{2}}^1\left(s-\frac{5}{6}\right)\frac{\partial f}{\partial s}\left(ta+(1-t)\,b,sc+(1-s)\,d\right)dsdt\Bigg\}.
\end{aligned}
$$

Computing these integrals and using the change of the variable $x=ta+(1-t)\,b$ and $y=sc+(1-s)\,d$ for $(t,s)\in[0,1]^2$, then multiplying both sides with $(b-a)(d-c)$, we get the desired result. □

Theorem 3. *Let $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ be a partial differentiable mapping on $\Delta=[a,b]\times[c,d]$. If $\frac{\partial^2 f}{\partial t\partial s}$ is a convex function on the co-ordinates on Δ, then the following inequality holds:*

$$
\begin{aligned}
& \Bigg|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9} \\
& +\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\Bigg| \\
\leq \quad & \frac{25\,(b-a)(d-c)}{72} \\
& \times\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|}{72}
\end{aligned}
$$

where

$$\begin{aligned} A &= \frac{1}{6(b-a)}\int_a^b \left[f(x,c)+4f\left(x,\frac{c+d}{2}\right)+f(x,d)\right]dx \\ &\quad +\frac{1}{6(d-c)}\int_c^d \left[f(a,y)+4f\left(\frac{a+b}{2},y\right)+f(b,y)\right]dy. \end{aligned}$$

Proof. By using Lemma 1, we can write

$$\begin{aligned} &\left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right. \\ &\left.+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\right| \\ \leq\ & (b-a)(d-c)\int_0^1\int_0^1 |p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}(ta+(1-t)b,sc+(1-s)d)\right|dtds. \end{aligned}$$

Since $f:\Delta\to\mathbb{R}$ is co-ordinated convex on Δ, we get

$$\begin{aligned} &\left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right. \\ &\left.+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\right| \\ \leq\ & (b-a)(d-c)\int_0^1 |q(y,s)|\left[\int_0^1 |p(x,t)|\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|\right.\right. \\ &\left.\left.+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|dt\right\}\right]ds. \end{aligned}$$

Computing the integral in the right hand side of above inequality, we have

$$\begin{aligned} &\int_0^1 |p(x,t)|\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|\right\}dt \\ =\ & \int_0^{\frac{1}{6}}\left(\frac{1}{6}-t\right)\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|\right\}dt \\ &+\int_{\frac{1}{6}}^{\frac{1}{2}}\left(t-\frac{1}{6}\right)\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|\right\}dt \\ &+\int_{\frac{1}{2}}^{\frac{5}{6}}\left(\frac{5}{6}-t\right)\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|\right\}dt \\ &+\int_{\frac{5}{6}}^{1}\left(t-\frac{5}{6}\right)\left\{t\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|+(1-t)\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|\right\}dt \end{aligned}$$

$$=\frac{5}{72}\left(\left|\frac{\partial^2 f}{\partial t\partial s}(a,sc+(1-s)d)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,sc+(1-s)d)\right|\right).$$

We obtain

$$\begin{aligned}
(2.2)\qquad & \left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right. \\
& +\frac{f\left(a,c\right)+f\left(b,c\right)+f\left(a,d\right)+f\left(b,d\right)}{36} \\
& \left.+\frac{1}{\left(b-a\right)\left(d-c\right)}\int_a^b\int_c^d f\left(x,y\right)dydx-A\right| \\
\leq\ & \frac{5\left(b-a\right)\left(d-c\right)}{72} \\
& \times\int_0^1\left|q\left(y,s\right)\right|\left\{\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,sc+\left(1-s\right)d\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,sc+\left(1-s\right)d\right)\right|\right\}ds
\end{aligned}$$

By a similar argument for the above integral, we have

$$\begin{aligned}
& \int_0^1\left|q\left(y,s\right)\right|\left\{\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,sc+\left(1-s\right)d\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,sc+\left(1-s\right)d\right)\right|\right\}ds \\
(2.3)\ =\ & \int_0^{\frac{1}{6}}\left(\frac{1}{6}-s\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,d\right)\right|\right\}ds \\
& +\int_0^{\frac{1}{6}}\left(\frac{1}{6}-s\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,d\right)\right|\right\}ds \\
& +\int_{\frac{1}{6}}^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,d\right)\right|\right\}ds \\
& +\int_{\frac{1}{6}}^{\frac{1}{2}}\left(s-\frac{1}{6}\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,d\right)\right|\right\}ds \\
& +\int_{\frac{1}{2}}^{\frac{5}{6}}\left(\frac{5}{6}-s\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,d\right)\right|\right\}ds \\
& +\int_{\frac{1}{2}}^{\frac{5}{6}}\left(\frac{5}{6}-s\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,d\right)\right|\right\}ds \\
& +\int_{\frac{5}{6}}^{1}\left(s-\frac{5}{6}\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,d\right)\right|\right\}ds \\
& +\int_{\frac{5}{6}}^{1}\left(s-\frac{5}{6}\right)\left\{s\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,c\right)\right|+\left(1-s\right)\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,d\right)\right|\right\}ds \\
=\ & \frac{5\left[\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,c\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,d\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,c\right)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,d\right)\right|\right]}{72}
\end{aligned}$$

If we use (2.3) in (2.2), we get the required result. $\square$

Theorem 4. *Let* $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta=[a,b]\times[c,d]$. *If* $\frac{\partial^2 f}{\partial t\partial s}$ *is bounded, i.e.,*

$$\left\|\frac{\partial^2 f}{\partial t\partial s}\left(ta+(1-t)b,sc+(1-s)d\right)\right\|_\infty=\sup_{(x,y)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f}{\partial t\partial s}\left(ta+(1-t)b,sc+(1-s)d\right)\right|<\infty$$

for all $(t,s)\in[0,1]^2$. *Then the following inequality holds:*

$$\begin{aligned}&\left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right.\\&+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}\\&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\right|\\\leq\;&\frac{25(b-a)(d-c)}{1296}\left\|\frac{\partial^2 f}{\partial t\partial s}(ta+(1-t)b,sc+(1-s)d)\right\|_\infty.\end{aligned}$$

Proof. From Lemma 1 and using the property of modulus, we have

$$\begin{aligned}&\left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right.\\&\left.+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\right|\\\leq\;&(b-a)(d-c)\int_0^1\int_0^1|p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}(ta+(1-t)\,b,sc+(1-s)\,d)\right|dtds.\end{aligned}$$

Since $\frac{\partial^2 f}{\partial t\partial s}$ is bounded, we have

$$\begin{aligned}&\left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right.\\&+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}\\&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\right|\\\leq\;&(b-a)(d-c)\\&\times\left\|\frac{\partial^2 f}{\partial t\partial s}(ta+(1-t)\,b,sc+(1-s)\,d)\right\|_\infty\int_0^1\int_0^1|p(x,t)\,q(y,s)|\,dtds.\end{aligned}\tag{2.4}$$

By a simple calculation,

$$\int_0^1\int_0^1|p(x,t)\,q(y,s)|\,dtds=\frac{25}{1296}.\tag{2.5}$$

If we use (2.5) in (2.4), we have

$$\begin{aligned}&\left|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\right.\\&\left.+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\right|\\\leq\;&\frac{25(b-a)(d-c)}{1296}\left\|\frac{\partial^2 f}{\partial t\partial s}(ta+(1-t)\,b,sc+(1-s)\,d)\right\|_\infty.\end{aligned}$$

This completes the proof. □

References

[1] Akdemir, A.O. and Özdemir M.E., Some Hadamard-type inequalities for co-ordinated $P-$convex functions and Godunova-Levin functions, American Institu of Physics Conference Proceedings, 2010 [ONLINE: http://scitation.aip.org/dbt/dbt.jsp?KEY=APCPCS&Volume=1309&Issue=1].

[2] Latif, M.A. and Alomari, M., On Hadamard-type inequalities for $h-$convex functions on the co-ordinates, International Journal of Math. Analysis, 3 (2009), no:33, 1645-1656.

[3] Latif, M.A. and Alomari, M., Hadamard-type inequalities for product two convex functions on the co-ordinates, International Mathematical Forum, 4 (2009),no:47, 2327-2338.

[4] Bakula, M.K. and Pecaric, J., On the Jensen's inequality for convex functions on the co-ordinates in a rectangle from the plane, Taiwanese Journal of Math., 5 (2006), 1271-1292.

[5] Alomari, M. and Darus, M., The Hadamard's inequality for $s-$convex function of $2-$variables on the co-ordinates, International Journal of Math. Analysis, 2 (2008) no:13, 629-638.

[6] Alomari, M. and Darus, M., Hadamard-type inequalities for $s-$convex functions, International Mathematical Forum, 3 (2008), no:40, 1965-1975.

[7] Alomari, M. and Darus, M., Co-ordinated $s-$convex function in the first sense with some Hadamard-type inequalities, Int. Journal Contemp. Math. Sciences, 3 (2008), no:32, 1557-1567.

[8] Hwang, D.Y., Tseng, K.L. and Yang, G.S., Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane, Taiwanese Journal of Mathematics, 11 (2007), 63-73.

[9] Özdemir, M.E., Set, E. and Sarıkaya, M.Z., Some new Hadamard's type inequalities for co-ordinated $m-$convex and $(\alpha, m)-$convex functions, Accepted.

[10] Sarıkaya, M.Z., Set, E., Özdemir, M.E. and Dragomir, S. S., New some Hadamard's type inequalities for co-ordinated convex functions, Accepted.

[11] Dragomir, S.S., On the Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, Taiwanese Journal of Mathematics, 5 (2001), no:4, 775-788.

[12] Sarıkaya, M.Z., Set, E. and Özdemir, M.E., On new inequalities of Simpson's type for convex functions, Submitted.

[13] Set, E., Özdemir, M.E. and Sarıkaya, M.Z., On new inequalities of Simpson's type for quasi-convex functions with applications, Submitted.

[14] Sarıkaya, M.Z., Set, E. and Özdemir, M.E., On new inequalities of Simpson's type for $s-$convex functions, Computers & Mathematics with Applications, 60, 8 (2010).

[15] Liu, B.Z., An inequality of Simpson type, Proc. R. Soc. A, 461 (2005), 2155-2158.

[16] Dragomir, S.S., Agarwal, R.P. and Cerone, P., On Simpson's inequality and applications, J. of Ineq. and Appl., 5 (2000), 533-579.

[17] Alomari, M., Darus, M. and Dragomir, S.S., New inequalities of Simpson's type for $s-$convex functions with applications, RGMIA Res. Rep. Coll., 12 (4) (2009), Article 9.

[18] Ujevic, N., Double integral inequalities of Simpson type and applications, J. Appl. Math. and Computing, 14 (2004), no:1-2, p. 213-223.

[19] Zhongxue, L., On sharp inequalities of Simpson type and Ostrowski type in two independent variables, Comp. and Math. with Appl., 56 (2008), 2043-2047.

[♦]Ataturk University, K.K. Education Faculty, Department of Mathematics, 25240, Erzurum, Turkey
E-mail address: `emos@atauni.edu.tr`

[♠]Ağri İbrahim Çeçen University, Faculty of Science and Arts, Department of Mathematics, 04100, Ağri, Turkey
E-mail address: `ahmetakdemir@agri.edu.tr`

[♦]Ataturk University, K.K. Education Faculty, Department of Mathematics, 25240, Erzurum, Turkey
E-mail address: `hkavurmaci@atauni.edu.tr`

E-mail address: `merveavci@ymail.com`